


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 211/56, 401/06, A61K 31/445	A1	(11) International Publication Number: WO 95/20575 (43) International Publication Date: 3 August 1995 (03.08.95)
(21) International Application Number: (22) International Filing Date: 26 January (30) Priority Data: 9401642.5 28 January 1994 (2) 9401639.1 28 January 1994 (2) <div style="text-align: center; font-size: 2em; margin-top: 20px;">77</div> (71) Applicant (for all designated States except GB/US): SHARP & DOHME LIMITED [GB/GB]; Hertford Road, Hoddesdon, Hertfordshire EN11 9BU (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): MACCOSS, Malcolm [GB/US]; 126 E. Lincoln Avenue, Rahway, NJ 07065-0900 (US). SWAIN, Christopher, John [GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). (74) Agent: HISCOCK, Ian, James; Merck & Co., Inc., Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB).		es: AM, AT, AU, BB, BG, BR, BY, CA, CH, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, LK, LR, LT, LU, LV, MD, MG, MN, MW, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, UZ, VN, European patent (AT, BE, CH, DE, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI, CF, CG, CI, CM, GA, GN, ML, MR, NE, ARIPO patent (KE, MW, SD, SZ). <i>wun international search report.</i>
(54) Title: ARALKYLAMINO SUBSTITUTED AZACYCLIC THERAPEUTIC AGENTS <div style="text-align: center;">  <p>(I)</p> </div> (57) Abstract <p>The present invention relates to compounds of formula (I) wherein n is 1, 2 or 3 and where any carbon atom of (CH₂)_n may be substituted by R⁴ and/or R⁵; R¹ represents (CH₂)_qphenyl, wherein q is zero, 1, 2 or 3, which may be optionally substituted in the phenyl ring; R² represents aryl selected from phenyl and naphthyl; heteroaryl selected from indazolyl, thienyl, furyl, pyridyl, thiazolyl, tetrazolyl and quinolyl; benzhydryl; or benzyl; wherein each aryl or heteroaryl moiety may be substituted; R³ represents H or C₁-alkyl; R⁴ and R⁵ each independently represent H, halo, C₁-alkyl, oxo, CO₂R^a or CONR^aR^b; R⁶ represents H or C₁-alkyl; R⁷ represents H, C₁-alkyl optionally substituted by a hydroxy group, or (CH₂)_pNR⁹R¹⁰, CO₂R¹⁶, CONR⁹R¹⁰, (CH₂)_pCO₂R¹⁶, (CH₂)_pCONR⁹R¹⁰, (CH₂)_pNR⁹COR¹⁶, (CH₂)_pNHSO₂R¹¹, (CH₂)_pOR¹⁶, (CH₂)_pOC(O)R⁹, (CH₂)_pOC₁₋₄alkylCOR¹⁷, or optionally substituted phenyl; R⁸ represents H, COR^a, CO₂R^a, COCONR^aR^b, COCO₂R^a, C₁-alkyl optionally substituted by a group selected from (CO₂R^a, CONR^aR^b, hydroxy, cyano, COR^a, NR^aR^b, C(NOH)NR^aR^b, CONHphenyl(C₁₋₄alkyl), COCO₂R^a, CONHNHNR^aR^b, C(S)NR^aR^b, CONR^aC₁₋₆alkylR¹², CONR¹³C₂₋₆alkynyl, CONR¹³C₂₋₆alkenyl, COCONR^aR^b, CONR^aC(NR^b)NR^aR^b, CONR^aheteroaryl, and optionally substituted phenyl) or C₁-alkyl, optionally substituted by oxo, substituted by an optionally substituted aromatic heterocycle; with the proviso that when R⁶ and R⁷ both represent H, R⁸ represents C₁-alkyl optionally substituted by oxo, substituted by an optionally substituted aromatic heterocycle selected from pyrrolyl, pyrazolyl, pyrazinyl, pyridazinyl, oxadiazolyl, thiadiazolyl, imidazolyl, benzimidazolyl, benzoxazolyl, benzthiophenyl, benzofuranyl and indolyl, or a substituted aromatic heterocycle selected from thienyl, furyl, pyridyl, triazolyl, tetrazolyl, thiazolyl, oxazolyl, quinolyl, isoxazolyl and isothiazolyl; and R^a and R^b each independently represent H, C₁-alkyl, trifluoromethyl or phenyl optionally substituted by C₁-alkyl, halo or trifluoromethyl. The compounds are of particular use in the treatment of pain, inflammation, migraine and emesis.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

ARALKYLAMINO SUBSTITUTED AZACYCLIC THERAPEUTIC AGENTS

This invention relates to a class of azacyclic compounds, which
5 are useful as tachykinin antagonists. More particularly, the compounds of
the invention comprise an azacyclic ring system substituted by an
aralkylamino moiety.

The tachykinins are a group of naturally occurring peptides
found widely distributed throughout mammalian tissues, both within the
10 central nervous system and in peripheral nervous and circulatory systems.

The tachykinins are distinguished by a conserved
carboxyl-terminal sequence:

Phe-X-Gly-Leu-Met-NH₂

At present, there are three known mammalian tachykinins
15 referred to as substance P, neurokinin A (NKA, substance K, neuromedin L)
and neurokinin B (NKB, neuromedin K) (for review see J.E. Maggio, *Peptides*
(1985) 6(suppl. 3), 237-242). The current nomenclature designates the three
tachykinin receptors mediating the biological actions of substance P, NKA
and NKB as the NK₁, NK₂ and NK₃ receptors, respectively.

20 Evidence for the usefulness of tachykinin receptor antagonists
in pain, headache, especially migraine, Alzheimer's disease, multiple
sclerosis, attenuation of morphine withdrawal, cardiovascular changes,
oedema, such as oedema caused by thermal injury, chronic inflammatory
diseases such as rheumatoid arthritis, asthma/bronchial hyperreactivity and
25 other respiratory diseases including allergic rhinitis, inflammatory diseases of
the gut including ulcerative colitis and Crohn's disease, ocular injury and
ocular inflammatory diseases, proliferative vitreoretinopathy, irritable bowel
syndrome and disorders of bladder function including cystitis and bladder
detruser hyper-reflexia is reviewed in "Tachykinin Receptors and Tachykinin

- 2 -

Receptor Antagonists", C.A. Maggi, R. Patacchini, P. Rovero and A. Giachetti, *J. Auton. Pharmacol.* (1993) 13, 23-93.

For instance, substance P is believed inter alia to be involved in the neurotransmission of pain sensations [Otsuka *et al*, "Role of Substance P as a Sensory Transmitter in Spinal Cord and Sympathetic Ganglia" in 1982 *Substance P in the Nervous System*, Ciba Foundation Symposium 91, 13-34 (published by Pitman) and Otsuka and Yanagisawa, "Does Substance P Act as a Pain Transmitter?" *TIPS* (1987) 8, 506-510], specifically in the transmission of pain in migraine (B.E.B. Sandberg *et al*, *J. Med Chem.* (1982) 25, 1009) and in arthritis [Levine *et al Science* (1984) 226, 547-549].

Tachykinins have also been implicated in gastrointestinal (GI) disorders and diseases of the GI tract such as inflammatory bowel disease [Mantyh *et al Neuroscience* (1988) 25(3), 817-37 and D. Regoli in "Trends in Cluster Headache" Ed. Sicuteri *et al* Elsevier Scientific Publishers, Amsterdam (1987) page 85]] and emesis [F. D. Tattersall *et al, Eur. J. Pharmacol.*, (1993) 250, R5-R6]. It is also hypothesised that there is a neurogenic mechanism for arthritis in which substance P may play a role [Kidd *et al* "A Neurogenic Mechanism for Symmetrical Arthritis" in *The Lancet*, 11 November 1989 and Grönblad *et al*, "Neuropeptides in Synovium of Patients with Rheumatoid Arthritis and Osteoarthritis" in *J. Rheumatol.* (1988) 15(12), 1807-10]. Therefore, substance P is believed to be involved in the inflammatory response in diseases such as rheumatoid arthritis and osteoarthritis, and fibrositis [O'Byrne *et al, Arthritis and Rheumatism* (1990) 33, 1023-8]. Other disease areas where tachykinin antagonists are believed to be useful are allergic conditions [Hamelet *et al, Can. J. Pharmacol. Physiol.* (1988) 66, 1361-7], immunoregulation [Lotz *et al, Science* (1988) 241, 1218-21 and Kimball *et al, J. Immunol.* (1988) 141(10), 3564-9] vasodilation, bronchospasm, reflex or neuronal control of the viscera [Mantyh *et al, PNAS* (1988) 85, 3235-9] and, possibly by arresting or slowing

- 3 -

β -amyloid-mediated neurodegenerative changes [Yankner *et al*, *Science* (1990) 250, 279-82] in senile dementia of the Alzheimer type, Alzheimer's disease and Down's Syndrome.

5 Tachykinin antagonists may also be useful in the treatment of small cell carcinomas, in particular small cell lung cancer (SCLC) [Langdon *et al*, *Cancer Research* (1992) 52, 4554-7].

Substance P may also play a role in demyelinating diseases such as multiple sclerosis and amyotrophic lateral sclerosis [J. Luber-Narod *et al*, poster *C.I.N.P. XVIIIth Congress*, 28th June-2nd July 1992], and in
10 disorders of bladder function such as bladder detrusor hyper-reflexia (*Lancet*, 16th May 1992, 1239).

It has furthermore been suggested that tachykinins have utility in the following disorders: depression, dysthymic disorders, chronic obstructive airways disease, hypersensitivity disorders such as poison ivy,
15 vasospastic diseases such as angina and Reynauld's disease, fibrosing and collagen diseases such as scleroderma and eosinophilic fascioliasis, reflex sympathetic dystrophy such as shoulder/hand syndrome, addiction disorders such as alcoholism, stress related somatic disorders, neuropathy, neuralgia, disorders related to immune enhancement or suppression such as systemic
20 lupus erythmatosus (European patent specification no. 0 436 334), ophthalmic disease such as conjunctivitis, vernal conjunctivitis, and the like, and cutaneous diseases such as contact dermatitis, atopic dermatitis, urticaria, and other eczematoid dermatitis (European patent specification no. 0 394 989).


25 In view of their metabolic instability, peptide derivatives are likely to be of limited utility as therapeutic agents. It is for this reason that non-peptide tachykinin antagonists are sought.

In essence, this invention provides a class of potent non-peptide tachykinin antagonists. By virtue of their non-peptide nature, the

- 4 -

compounds of the present invention do not suffer from the shortcomings, in terms of metabolic instability, of the known peptide-based tachykinin antagonists discussed above.

The present invention provides a compound of formula (I), or a
 5 pharmaceutically acceptable salt or prodrug thereof:



wherein

10 n is 1, 2 or 3 and where any carbon atom of $(CH_2)_n$ may be substituted by R^4 and/or R^5 ;

R^1 represents $(CH_2)_q$ phenyl, wherein q is zero, 1, 2 or 3, which may be optionally substituted in the phenyl ring by 1, 2 or 3 groups selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halo, cyano, nitro, trifluoromethyl,
 15 trimethylsilyl, $-OR^a$, $-SR^a$, $-SOR^a$, $-SO_2R^a$, $-NR^aR^b$, $-NR^aCOR^b$, $-NR^aCO_2R^b$, $-COR^a$, $-CO_2R^a$ and $-CONR^aR^b$;

R^2 represents aryl selected from phenyl and naphthyl; heteroaryl selected from indazolyl, thienyl, furyl, pyridyl, thiazolyl, tetrazolyl and quinolyl; benzhydryl; or benzyl; wherein each aryl or heteroaryl moiety
 20 may be substituted by C_{1-6} alkyl, C_{1-6} alkoxy, halo or trifluoromethyl;

R^3 represents H or C_{1-6} alkyl;

R^4 and R^5 each independently represent H, halo, C_{1-6} alkyl, oxo, CO_2R^a or $CONR^aR^b$;

R^6 represents H or C_{1-6} alkyl;

- 5 -

R⁷ represents H, C₁₋₆alkyl optionally substituted by a hydroxy group, or (CH₂)_pNR⁹R¹⁰, CO₂R¹⁶, CONR⁹R¹⁰, (CH₂)_pCO₂R¹⁶, (CH₂)_pCONR⁹R¹⁰, (CH₂)_pNR⁹COR¹⁶, (CH₂)_pNHSO₂R¹¹, (CH₂)_pOR¹⁶, (CH₂)_pOC(O)R⁹, (CH₂)_pOC₁₋₄alkylCOR¹⁷, or phenyl optionally substituted by 1, 2 or 3 groups selected from C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, halo, cyano, nitro, trifluoromethyl, trimethylsilyl, -OR^a, -SR^a, -SOR^a, -SO₂R^a, -NR^aR^b, -NR^aCOR^b, -NR^aCO₂R^b, -COR^a, -CO₂R^a and -CONR^aR^b;

R⁸ represents H, COR^a, CO₂R^a, COCONR^aR^b, COCO₂R^a, C₁₋₆alkyl optionally substituted by a group selected from (CO₂R^a, CONR^aR^b, hydroxy, cyano, COR^a, NR^aR^b, C(NOH)NR^aR^b, CONHphenyl(C₁₋₄alkyl), COCO₂R^a, CONHNR^aR^b, C(S)NR^aR^b, CONR^aC₁₋₆alkylR¹², CONR¹³C₂₋₆alkynyl, CONR¹³C₂₋₆alkenyl, COCONR^aR^b, CONR^aC(NR^b)NR^aR^b, CONR^aheteroaryl, and phenyl optionally substituted by one or more substituents selected from C₁₋₆alkyl, C₁₋₆alkoxy, halo and trifluoromethyl) or C₁₋₆alkyl, optionally substituted by oxo, substituted by an optionally substituted aromatic heterocycle;

with the proviso that when R⁶ and R⁷ both represent H, R⁸ represents C₁₋₆alkyl optionally substituted by oxo, substituted by an optionally substituted aromatic heterocycle selected from pyrrolyl, pyrazolyl, pyrazinyl, pyridazinyl, oxadiazolyl, thiadiazolyl, imidazolyl, benzimidazolyl, benzoxazolyl, benzthiophenyl, benzofuranyl and indolyl, or a substituted aromatic heterocycle selected from thienyl, furyl, pyridyl, triazolyl, tetrazolyl, thiazolyl, oxazolyl, quinolyl, isoxazolyl and isothiazolyl;

R^a and R^b each independently represent H, C₁₋₆alkyl, trifluoromethyl or phenyl optionally substituted by C₁₋₆alkyl, halo or trifluoromethyl;

R⁹ and R¹⁰ each independently represent H or C₁₋₆alkyl;

R¹¹ represents NR¹⁴R¹⁵ or an optionally substituted aromatic or non-aromatic azacyclic or azabicyclic group;

- 6 -

R¹² represents OR^a, CONR^aR^b or heteroaryl;

R¹³ represents H or C₁₋₆alkyl;

R¹⁴ and R¹⁵ each independently represent H, C₁₋₆alkyl, phenyl optionally substituted by one or more of C₁₋₆alkyl, C₁₋₆alkoxy, halo or trifluoromethyl, or phenylC₁₋₄alkyl optionally substituted in the phenyl ring
5 or by one or more of C₁₋₆alkyl, C₁₋₆alkoxy, halo or trifluoromethyl;

R¹⁶ represents C₁₋₆alkyl;

R¹⁷ represents C₁₋₆alkoxy, amino, C₁₋₄alkylamino or di(C₁₋₄alkyl)amino; and

10 p is 1 to 4.

As used herein, the definition of each expression, when it occurs more than once in any structure, is intended to be independent of its definition elsewhere in the same structure.

The alkyl, alkenyl and alkynyl groups referred to with respect to
15 the above formula may represent straight, branched or cyclic groups. Thus, for example, suitable alkyl groups include methyl, ethyl, n- or iso-propyl, n-, sec-, iso- or tert-butyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, and cycloalkyl-alkyl groups such as cyclopropylmethyl; suitable alkenyl groups include vinyl and allyl; and suitable alkynyl groups include propargyl.

20 The term "halo" as used herein includes fluoro, chloro, bromo and iodo, especially chloro and fluoro.

The present invention includes within its scope prodrugs of the compounds of formula (I) above. In general, such prodrugs will be functional derivatives of the compounds of formula (I) which are readily convertible in vivo
25 into the required compound of formula (I). Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985.

The compounds according to the invention may exist both as enantiomers and as diastereomers. In particular, the relative orientation of

- 7 -

the 2- and 3- substituents on the azacyclic ring may give rise to cis and trans diastereoisomers, of which the cis stereochemistry is preferred. It is to be understood that all such isomers and mixtures thereof are encompassed within the scope of the present invention.

5 Preferably n is 2 or 3, more preferably 3.

Preferably q is zero and R¹ represents substituted phenyl.

When R¹ is substituted phenyl suitable substituents include nitro, trifluoromethyl, trimethylsilyl, bromo, chloro, fluoro, iodo, cyano, methyl, ethyl, cyclopropyl, vinyl, methoxy, phenoxy, amino and carbonylmethoxy.

10 Preferably R¹ represents phenyl substituted by one or more groups selected from methyl, trifluoromethyl, chloro and t-butyl.

Preferably R¹ represents disubstituted phenyl, more preferably 3,5-disubstituted phenyl such as 3,5-dichlorophenyl or 3,5-bis(trifluoromethyl)phenyl, or monosubstituted phenyl, such as

15 3-substituted phenyl, e.g. 3-t-butylphenyl.

Preferably R² represents unsubstituted benzhydryl, phenyl substituted by halo such as fluoro, for example 4-fluorophenyl, or unsubstituted phenyl, more preferably unsubstituted phenyl.

Preferably R⁴ and R⁵ both represent H.

20 Suitable values for R⁶ include H, methyl and ethyl. Preferably R⁶ represents H or methyl, more preferably H.

Preferably R⁷ represents C₁₋₆alkyl optionally substituted by a hydroxy group, such as methyl, ethyl, CH₂OH, CH₂CH₂OH, CH(OH)CH₃ or C(OH)(CH₃)₂, more preferably methyl or CH₂OH.

25 When R⁸ represents C₁₋₆alkyl, optionally substituted by oxo, substituted by a substituted aromatic heterocycle, suitable substituents in the heterocyclic ring include C₁₋₆alkyl, C₁₋₆alkoxy, oxo, thioxo, halo, trifluoromethyl, NR^aR^b, NR^aCOR^b, CONR^aR^b, CO₂R^a, SO₂R^a and CH₂OR^a, where R^a and R^b are as previously defined.

- 8 -

Preferably R^8 represents C_{1-3} alkyl such as methyl, ethyl or i-propyl substituted by a substituted or unsubstituted aromatic heterocycle.

Suitable heterocycles include thienyl, furyl, pyrrolyl, pyridyl, pyrazolyl, triazolyl, tetrazolyl, thiazolyl, pyrazinyl, pyridazinyl, oxazolyl, oxadiazolyl, thiadiazolyl, isoxazolyl, quinolyl, isothiazolyl, imidazolyl, benzimidazolyl, benzoxazolyl, benzothiophenyl, benzofuranyl and indolyl.

In one group of compounds according to the invention R^8 represents CH_2 -Het, $CH(CH_3)$ -Het, $C(CH_3)_2$ -Het or $C(O)$ -Het, where Het is pyrrolyl, pyrazolyl, pyrazinyl, pyridazinyl, oxadiazolyl, thiadiazolyl, imidazolyl, benzimidazolyl, benzoxazolyl, benzothiophenyl, benzofuranyl or indolyl.

Preferably R^8 represents CH_2 -Het, $CH(CH_3)$ -Het, $C(CH_3)_2$ -Het or $C(O)$ -Het where Het is substituted or unsubstituted oxazolyl, oxadiazolyl, tetrazolyl, thiazolyl, thiadiazolyl, furanyl, thienyl, triazolyl, pyrazinyl, pyridyl, pyridazinyl, imidazolyl or benzimidazolyl. More preferably Het is triazolyl or triazolyl substituted by oxo.

Other suitable values for R^8 include H, COR^a , CO_2R^a , $COCONR^aR^b$, $COCO_2R^a$, C_{1-6} alkyl and C_{1-6} alkyl substituted by a group selected from CO_2R^a , $CONR^aR^b$, CN, $C(NO)NR^aR^b$, $CONHphenyl(C_{1-4}alkyl)$, optionally substituted phenyl, $CONHNR^aR^b$, $COCONR^aR^b$, $CONR^aC(NH)NH_2$, $CSNR^aR^b$, $CONR^{13}C_{2-6}alkynyl$, $CONR^aC_{1-6}alkylR^{12}$ and $CONR^aheteroaryl$.

It will be appreciated that, when R^8 comprises a heteroaryl moiety substituted by an oxo or thioxo substituent, different tautomeric forms are possible so that the substituent on the heteroaryl moiety may be represented as =O or -OH, or =S or -SH, respectively. For the avoidance of doubt, all such tautomeric forms are embraced by the present invention.

When R^{11} represents $NR^{14}R^{15}$, R^{14} and R^{15} are preferably both C_{1-6} alkyl such as methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl or t-butyl. More preferably R^{14} and R^{15} will both represent methyl.

- 9 -

When R^{11} represents an aromatic or non-aromatic azacycle or azabicyclic it may contain one or more additional heteroatoms selected from O, S and N or groups NR^{18} , where R^{18} is H, C_{1-6} alkyl or phenyl C_{1-4} alkyl, and may be unsubstituted or substituted. Suitable substituents include

5 C_{1-6} alkyl, C_{1-6} alkoxy, oxo, SH, =S, halo, trifluoromethyl, NR^aR^b , NR^aCOR^b , $CONR^aR^b$, CO_2R^a and CH_2OR^a , where R^a and R^b are as previously defined.


When R^{11} represents an aromatic azacycle or azabicyclic, suitable values of R^{11} include imidazolyl, triazolyl, tetrazolyl, oxazolyl,

10 thiazolyl, pyrrolyl, pyrazolyl, pyrazinyl, pyridyl, oxadiazolyl, thiadiazolyl, isoxazolyl, isothiazolyl, benzimidazolyl, benzoxazolyl and indolyl, preferably imidazolyl, such as 2,4-imidazolyl, or pyridyl, more preferably pyridyl such as 4-, 3- or 2-pyridyl.

When R^{11} represents a non-aromatic azacycle or azabicyclic, suitable values of R^{11} include morpholinyl, piperdiny, pyrrolidinyl,

15 piperazinyl, methylpiperazinyl, azanorbornanyl, azabicyclo[2.2.2]octanyl and azabicyclo[3.2.2]nonyl, preferably morpholinyl, pyrrolidinyl, methylpiperazinyl, quinuclidinyl (azabicyclo[2.2.2] octanyl) or azabicyclo[3.2.2]nonyl, more preferably pyrrolidinyl.

20 A particular sub-class of compounds according to the present invention is represented by compounds of formula (Ia), and pharmaceutically acceptable salts and prodrugs thereof:



- 10 -


wherein

R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^8 and n are as defined for formula (I) above; and

5 R^7 represents C_{1-6} alkyl optionally substituted by a hydroxy group, or $(CH_2)_pNR^9R^{10}$, CO_2R^{16} , $CONR^9R^{10}$, $(CH_2)_pCO_2R^{16}$, $(CH_2)_pCONR^9R^{10}$, $(CH_2)_pNR^9COR^{16}$, $(CH_2)_pNHSO_2R^{11}$, $(CH_2)_pOR^{16}$, $(CH_2)_pOC(O)R^9$, $(CH_2)_pOC_{1-4}alkylCOR^{17}$, or phenyl optionally substituted by 1, 2 or 3 groups selected from C_{1-6} alkyl, $C_{2-6}alkenyl$, $C_{2-6}alkynyl$, halo, cyano, nitro, trifluoromethyl, trimethylsilyl, $-OR^a$, $-SR^a$, $-SOR^a$, $-SO_2R^a$, $-NR^aR^b$, $-NR^aCOR^b$, $-NR^aCO_2R^b$, $-COR^a$, $-CO_2R^a$ and $-CONR^aR^b$.

A preferred sub-class of compounds according to the present invention is represented by compounds of formula (Ib), and pharmaceutically acceptable salts and prodrugs thereof:

15



wherein

20 R^2 , R^3 , R^4 , R^5 , R^6 , R^8 and n are as defined for formula (I) above;


R^{18} represents C_{1-6} alkyl optionally substituted by a hydroxy group; and

R^{19} represents phenyl optionally substituted by 1, 2 or 3 groups selected from C_{1-6} alkyl, $C_{2-6}alkenyl$, $C_{2-6}alkynyl$, halo, cyano, nitro,

- 11 -

trifluoromethyl, trimethylsilyl, $-OR^a$, $-SR^a$, $-SOR^a$, $-SO_2R^a$, $-NR^aR^b$, $-NR^aCOR^b$, $-NR^aCO_2R^b$, $-COR^a$, $-CO_2R^a$ and $-CONR^aR^b$.

A further preferred sub-class of compounds according to the present invention is represented by compounds of formula (Ic) and pharmaceutically acceptable salts and prodrugs thereof:




wherein

- 10 R^3 and R^8 are as defined for formula (I);
 R^{20} and R^{21} independently represent H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, chloro, bromo, fluoro, iodo, cyano, nitro, trifluoromethyl, trimethylsilyl, OR^a , SR^a , SOR^a , SO_2R^a , NR^aR^b , NR^aCOR^b , $NR^aCO_2R^b$, COR^a , CO_2R^a or $CONR^aR^b$, where R^a and R^b are as
15 previously defined;
 R^{22} is methyl optionally substituted by hydroxy;
 R^{23} is H or methyl; and
 R^{24} represents phenyl or benzhydryl wherein any of the phenyl rings of the phenyl or benzhydryl moieties may optionally be substituted by
20 halo or trifluoromethyl, preferably unsubstituted phenyl.
Particular values of R^{20} and R^{21} include methyl, ethyl, t-butyl, chloro, fluoro and trifluoromethyl. Preferably R^{20} and R^{21} are both other than hydrogen and are located at the 3- and 5-positions of the phenyl ring.

- 12 -

A preferred group of compounds according to the present invention are compounds of formula (Ic) wherein R^8 is optionally substituted triazolyl.

Another particular sub-class of compounds according to the present invention is represented by compounds of formula (Id), and pharmaceutically acceptable salts and prodrugs thereof:




wherein

R^1 , R^2 , R^3 , R^4 , R^5 , and n are as defined for formula (I) above; and

R^8 represents C_{1-6} alkyl, optionally substituted by oxo, substituted by an optionally substituted aromatic heterocycle selected from pyrrolyl, pyrazolyl, pyrazinyl, pyridazinyl, oxadiazolyl, thiadiazolyl, imidazolyl, benzimidazolyl, benzoxazolyl, benzthiophenyl, benzofuranyl and indolyl, or a substituted aromatic heterocycle selected from thienyl, furyl, pyridyl, triazolyl, tetrazolyl, thiazolyl, oxazolyl, quinolyl, isoxazolyl and isothiazolyl.

Another particularly preferred sub-class of compounds according to the present invention is represented by compounds of formula (Ie), and pharmaceutically acceptable salts and prodrugs thereof:

- 13 -




wherein R², R³, R⁴, R⁵, R⁸ and n are as defined for formula (Id) above; and

R¹⁹ represents phenyl optionally substituted by 1, 2 or 3

5 groups selected from C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, halo, cyano, nitro, trifluoromethyl, trimethylsilyl, -OR^a, -SR^a, -SOR^a, -SO₂R^a, -NR^aR^b, -NR^aCOR^b, -NR^aCO₂R^b, -COR^a, -CO₂R^a and -CONR^aR^b.

A yet further preferred sub-class of compounds according to the present invention is represented by compounds of formula (If) and pharmaceutically acceptable salts and prodrugs thereof:



wherein

15 R^3 and R^8 are as defined for formula (I);

R²⁰ and R²¹ independently represent H, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, chloro, bromo, fluoro, iodo, cyano, nitro, trifluoromethyl, trimethylsilyl, OR^a, SR^a, SOR^a, SO₂R^a, NR^aR^b, NR^aCOR^b,

- 14 -

$\text{NR}^a\text{CO}_2\text{R}^b$, COR^a , CO_2R^a or CONR^aR^b , where R^a and R^b are as previously defined; and

R^{24} represents phenyl or benzhydryl wherein any of the phenyl rings of the phenyl or benzhydryl moieties may optionally be substituted by halo or trifluoromethyl, preferably unsubstituted phenyl.

Particular values of R^{20} and R^{21} include methyl, ethyl, t-butyl, chloro, fluoro and trifluoromethyl. Preferably R^{20} and R^{21} are both other than hydrogen and are located at the 3- and 5-positions of the phenyl ring.

A preferred group of compounds according to the invention are compounds of formula (If) wherein R^8 is substituted triazolyl.

Specific compounds within the scope of the present invention include:

cis-3-((3,5-bis(trifluoromethyl))(α -methyl)benzylamino)-2-(4-fluorophenyl) piperidine;
cis-[3-((3,5-bis(trifluoromethyl))benzylamino)-2-(4-fluorophényl)piperidine-1-ylmethyl]-2,4-dihydro-[1,2,4]-triazol-3-one;
and pharmaceutically acceptable salts and prodrugs thereof.

For use in medicine, the salts of the compounds of formula (I) will be pharmaceutically acceptable salts. Other salts (such as the dibenzoyltartrate salts) may, however, be useful in the preparation of the compounds according to the invention or of their pharmaceutically acceptable salts. Suitable pharmaceutically acceptable salts of the compounds of this invention include acid addition salts which may, for example, be formed by mixing a solution of the compound according to the invention with a solution of a pharmaceutically acceptable acid such as hydrochloric acid, sulphuric acid, oxalic acid, fumaric acid, maleic acid, succinic acid, acetic acid, citric acid, tartaric acid, carbonic acid, phosphoric acid or p-toluenesulphonic acid. Salts of amine groups may also comprise quaternary ammonium salts in which the amino nitrogen atom carries a

- 15 -

suitable organic group such as an alkyl, alkenyl, alkynyl or aralkyl moiety. Furthermore, where the compounds of the invention carry an acidic moiety, suitable pharmaceutically acceptable salts thereof may include metal salts such as alkali metal salts, e.g. sodium or potassium salts; and alkaline earth
5 metal salts, e.g. calcium or magnesium salts.

Preferred salts of the compounds according to the invention include the hydrochloride and p-toluenesulphonic acid salts.

The invention also provides pharmaceutical compositions comprising a compound of this invention in association with a
10 pharmaceutically acceptable carrier. Preferably these compositions are in unit dosage forms such as tablets, pills, capsules, powders, granules, solutions or suspensions, or suppositories, for oral, parenteral or rectal administration, or administration by inhalation or insufflation.

For preparing solid compositions such as tablets, the principal
15 active ingredient is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums, and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the
20 present invention, or a non-toxic pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules. This solid
25 preformulation composition is then subdivided into unit dosage forms of the type described above containing from 0.1 to about 500 mg of the active ingredient of the present invention. The tablets or pills of the novel composition can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or

- 16 -

pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions, and flavoured emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinylpyrrolidone or gelatin.

Compositions for inhalation or insufflation include solutions and suspensions in pharmaceutically acceptable, aqueous or organic solvents, or mixtures thereof, and powders. The liquid or solid compositions may contain suitable pharmaceutically acceptable excipients as set out above. Preferably the compositions are administered by the oral or nasal respiratory route for local or systemic effect. Compositions in preferably sterile pharmaceutically acceptable solvents may be nebulised by use of inert gases. Nebulised solutions may be breathed directly from the nebulising device or the nebulising device may be attached to a face mask, tent or intermittent positive pressure breathing machine. Solution, suspension or powder compositions may be administered, preferably orally or nasally, from devices which deliver the formulation in an appropriate manner.

- 17 -

The compounds of formula (I) are of value in the treatment of a wide variety of clinical conditions which are characterised by the presence of an excess of tachykinin, in particular substance P, activity. These may include disorders of the central nervous system such as anxiety, depression, psychosis and schizophrenia; epilepsy; neurodegenerative disorders such as dementia, including senile dementia of the Alzheimer type, Alzheimer's disease and Down's syndrome; demyelinating diseases such as MS and ALS and other neuropathological disorders such as peripheral neuropathy, for example diabetic and chemotherapy-induced neuropathy, and postherpetic and other neuralgias; small cell carcinomas such as small cell lung cancer; respiratory diseases, particularly those associated with excess mucus secretion such as chronic obstructive airways disease, bronchopneumonia, chronic bronchitis, cystic fibrosis and asthma, and bronchospasm; inflammatory diseases such as inflammatory bowel disease, psoriasis, fibrositis, osteoarthritis, rheumatoid arthritis, pruritis and sunburn; allergies such as eczema and rhinitis; hypersensitivity disorders such as poison ivy; ophthalmic diseases such as conjunctivitis, vernal conjunctivitis, and the like; ophthalmic conditions associated with cell proliferation such as proliferative vitreoretinopathy; cutaneous diseases such as contact dermatitis, atopic dermatitis, urticaria, and other eczematoid dermatitis; addiction disorders such as alcoholism; stress related somatic disorders; reflex sympathetic dystrophy such as shoulder/hand syndrome; dysthymic disorders; adverse immunological reactions such as rejection of transplanted tissues and disorders related to immune enhancement or suppression such as systemic lupus erythematosus; gastrointestinal (GI) disorders and diseases of the GI tract such as disorders associated with the neuronal control of viscera, ulcerative colitis, Crohn's disease, irritable bowel syndrome and emesis, including acute, delayed or anticipatory emesis such as emesis induced by chemotherapy, radiation, toxins, viral or bacterial infections, pregnancy,

- 18 -

vestibular disorders, motion, surgery, migraine, and variations in intercranial pressure, in particular, for example, drug or radiation induced emesis or post-operative nausea and vomiting; disorders of bladder function such as cystitis, bladder detrusor hyper-reflexia and incontinence; fibrosing and collagen diseases such as scleroderma and eosinophilic fascioliasis; disorders of blood flow caused by vasodilation and vasospastic diseases such as angina, migraine and Reynaud's disease; and pain or nociception, for example, that attributable to or associated with any of the foregoing conditions, especially the transmission of pain in migraine.

The compounds of formula (I) are also of value in the treatment of a combination of the above conditions, in particular in the treatment of combined post-operative pain and post-operative nausea and vomiting.

The compounds of formula (I) are particularly useful in the treatment of emesis, including acute, delayed or anticipatory emesis, such as emesis induced by chemotherapy, radiation, toxins, pregnancy, vestibular disorders, motion, surgery, migraine, and variations in intercranial pressure. Most especially, the compounds of formula (I) are of use in the treatment of emesis induced by antineoplastic (cytotoxic) agents including those routinely used in cancer chemotherapy.

Examples of such chemotherapeutic agents include alkylating agents, for example, nitrogen mustards, ethyleneimine compounds, alkyl sulphonates and other compounds with an alkylating action such as nitrosoureas, cisplatin and dacarbazine; antimetabolites, for example, folic acid, purine or pyrimidine antagonists; mitotic inhibitors, for example, vinca alkaloids and derivatives of podophyllotoxin; and cytotoxic antibiotics.

Particular examples of chemotherapeutic agents are described, for instance, by D. J. Stewart in *Nausea and Vomiting: Recent Research and Clinical Advances*, Eds. J. Kucharczyk et al, CRC Press Inc., Boca Raton, Florida, USA (1991) pages 177-203, especially page 188. Commonly used

- 19 -

chemotherapeutic agents include cisplatin, dacarbazine (DTIC), dactinomycin, mechlorethamine (nitrogen mustard), streptozocin, cyclophosphamide, carmustine (BCNU), lomustine (CCNU), doxorubicin (adriamycin), daunorubicin, procarbazine, mitomycin, cytarabine, etoposide, methotrexate, 5-fluorouracil, vinblastine, vincristine, bleomycin and chlorambucil [R. J. Gralla *et al* in *Cancer Treatment Reports* (1984) 68(1), 163-172].

The compounds of formula (I) are also of use in the treatment of emesis induced by radiation including radiation therapy such as in the treatment of cancer, or radiation sickness; and in the treatment of post-operative nausea and vomiting.

It will be appreciated that the compounds of formula (I) may be presented together with another therapeutic agent as a combined preparation for simultaneous, separate or sequential use for the relief of emesis. Such combined preparations may be, for example, in the form of a twin pack.

A further aspect of the present invention comprises the compounds of formula (I) in combination with a 5-HT₃ antagonist, such as ondansetron, granisetron or tropisetron, or other anti-emetic medicaments, for example, a dopamine antagonist such as metoclopramide. Additionally, a compound of formula (I) may be administered in combination with an anti-inflammatory corticosteroid, such as dexamethasone. Furthermore, a compound of formula (I) may be administered in combination with a chemotherapeutic agent such as an alkylating agent, antimetabolite, mitotic inhibitor or cytotoxic antibiotic, as described above. In general, the currently available dosage forms of the known therapeutic agents for use in such combinations will be suitable.

When tested in the ferret model of cisplatin-induced emesis described by F. D. Tattersall *et al*, in *Eur. J. Pharmacol.*, (1993) 250, R5-R6,

- 20 -

the compounds of the present invention were found to attenuate the retching and vomiting induced by cisplatin.

The compounds of formula (I) are also particularly useful in the treatment of pain or nociception and/or inflammation and disorders
5 associated therewith such as, for example, neuropathy, such as diabetic and chemotherapy-induced neuropathy, postherpetic and other neuralgias, asthma, osteoarthritis, rheumatoid arthritis and especially migraine.

The present invention further provides a compound of formula (I) for use in therapy.

10 According to a further or alternative aspect, the present invention provides a compound of formula (I) for use in the manufacture of a medicament for the treatment of physiological disorders associated with an excess of tachykinins, especially substance P.

The present invention also provides a method for the the
15 treatment or prevention of physiological disorders associated with an excess of tachykinins, especially substance P, which method comprises administration to a patient in need thereof of a tachykinin reducing amount of a compound of formula (I) or a composition comprising a compound of formula (I).

20 For the treatment of certain conditions it may be desirable to employ a compound according to the present invention in conjunction with another pharmacologically active agent. For example, for the treatment of respiratory diseases such as asthma, a compound of formula (I) may be used in conjunction with a bronchodilator, such as a β_2 -adrenergic receptor
25 antagonist or tachykinin antagonist which acts at NK-2 receptors. The compound of formula (I) and the bronchodilator may be administered to a patient simultaneously, sequentially or in combination.

The present invention accordingly provides a method for the treatment of a respiratory disease, such as asthma, which method comprises

- 21 -

administration to a patient in need thereof of an effective amount of a compound of formula (I) and an effective amount of a bronchodilator.

The present invention also provides a composition comprising a compound of formula (I), a bronchodilator, and a pharmaceutically acceptable carrier.

The excellent pharmacological profile of the compounds of the present invention offers the opportunity for their use in therapy at low doses thereby minimising the risk of unwanted side effects.


In the treatment of the conditions associated with an excess of tachykinins, a suitable dosage level is about 0.001 to 50 mg/kg per day, in particular about 0.01 to about 25 mg/kg, such as from about 0.05 to about 10 mg/kg per day.

For example, in the treatment of conditions involving the neurotransmission of pain sensations, a suitable dosage level is about 0.001 to 25 mg/kg per day, preferably about 0.005 to 10 mg/kg per day, and especially about 0.005 to 5 mg/kg per day. The compounds may be administered on a regimen of 1 to 4 times per day, preferably once or twice per day.

In the treatment of emesis using an injectable formulation, a suitable dosage level is about 0.001 to 10 mg/kg per day, preferably about 0.005 to 5 mg/kg per day, and especially 0.01 to 2 mg/kg per day. The compounds may be administered on a regimen of 1 to 4 times per day, preferably once or twice per day.

It will be appreciated that the amount of a compound of formula (I) required for use in any treatment will vary not only with the particular compounds or composition selected but also with the route of administration, the nature of the condition being treated, and the age and condition of the patient, and will ultimately be at the discretion of the attendant physician.

The compounds according to the present invention may be prepared by a process (A) which comprises reacting a compound of formula (II) with a compound of formula (III):



wherein R¹, R², R⁴, R⁵, R⁶, R⁷, R⁸ and n are as defined for formula (I), except that any reactive moiety is protected by a suitable protecting group; and one of R³⁰ and R³¹ represents a leaving group and the other of R³⁰ and R³¹ represents NHR³, where R³ is as defined for formula (I); in the presence of a base, followed by deprotection, if required.

Suitably R³⁰ represents NHR³ and R³¹ represents a leaving group.

Suitable leaving groups include halo, e.g. chloro, bromo or iodo, or sulphonate derivatives such as tosylate, mesylate or triflate.

The reaction is conveniently carried out in a suitable organic solvent, such as an ether, e.g. 1,2-dimethoxyethane, at a temperature in the region of 0°C. Favoured bases of use in the reaction include alkali metal amides and hydrides, such as potassium bis(trimethylsilyl)amide or potassium hydride. Suitably, sodium hydride is used.

According to another process (B), compounds of formula (I) wherein R⁶ is hydrogen, may be prepared by the reductive amination of a compound of formula (II) in which R³⁰ is the group NHR³, with a compound of formula R¹COR⁷, in the presence of a reducing agent. Suitable reducing

- 23 -

agents for use in this reaction include, for example, sodium
cyanoborohydride or sodium triacetoxyborohydride, or catalytic
hydrogenation. The reaction is conveniently effected in a suitable solvent
such as acetic acid or methanol at a temperature between 0°C and 50°C,
5 conveniently at about room temperature.

Alternatively, according to a further process (C), compounds of
formula (I) may be prepared from different compounds of formula (I) by
interconversion processes. In particular, interconversion processes may be
used to vary the group R^8 . For example, compounds of formula (I) wherein
10 R^8 is other than H may be prepared from the corresponding compounds of
formula (I) wherein R^8 is H by reaction with a reagent suitable to introduce
the group R^8 , for example, a halide or acyl halide, or corresponding mesylate
or tosylate, of formula R^8-L , where L represents halo, such as chloro, bromo
or iodo, methylsulphonate or p- toluenesulphonate, or any other suitable
15 leaving group, in the presence of a base. Suitable bases of use in the
reaction include inorganic bases such as alkali metal carbonates, for
example, potassium carbonate. Conveniently the reaction is effected in a
suitable organic solvent, for example, dimethylformamide.

Compounds of formula (I) wherein R^8 is COR^a may be
20 prepared from compounds of formula (I) wherein R^8 is H by, for example,
reaction with an appropriate acid anhydride.


Compounds of formula (I) wherein R^8 is C_{1-6} alkyl may be
prepared from corresponding compounds of formula (I) wherein R^8 is COR^a
by reduction using, for example, borane or a borohydride such as sodium
25 cyanoborohydride.

Compounds of formula (I) wherein R^8 is C_{1-6} alkyl substituted
by $CONR^aR^b$ may be prepared from corresponding compounds of formula (I)
wherein R^8 is C_{1-6} alkyl substituted by CO_2R^a by treatment with ammonia or
an amine of formula NR^aR^b .

- 24 -

Compounds of formula (I) wherein R⁸ is C₁₋₆alkyl substituted by 5-oxadiazolyl may be prepared from compounds of formula (I) wherein R⁸ is C₁₋₆alkyl substituted by CO₂R^a, where R^a represents C₁₋₆alkyl, by reaction with a compound of formula (IV)

5



wherein R³² represents H or a suitable substituent, in the presence of a base.

10

Suitable bases of use in the reaction include alkali metals, such as, for example, sodium, and alkali metal hydrides, such as, for example, sodium hydride.

The reaction is conveniently effected in a suitable organic solvent. Which solvents will be appropriate will depend on the nature of the base used. For example, where the base used is an alkali metal, suitable solvents will include alcohols, for example, ethanol, whereas where the base used is an alkali hydride, suitable solvents will include ethers, for example, tetrahydrofuran.

15

Preferably the reaction is conducted at elevated temperature, such as the reflux temperature of the chosen solvent.

20

Compounds of formula (I) wherein R⁸ is C₁₋₆alkyl substituted by tetrazolyl may be prepared from compounds of formula (I) wherein R⁸ is C₁₋₆alkyl substituted by cyano by treatment with an alkali metal azide, such as sodium azide.

25

Compounds of formula (I) wherein R⁸ is C₁₋₆alkyl substituted by thiazolyl may be prepared from compounds of formula (I) wherein R⁸ is C₁₋₆alkyl substituted by CSNH₂ by reaction with a compound of formula


- 25 -

Hal-CH₂C(O)-R⁶⁰, where Hal is halo, such as bromo, chloro or iodo, and R⁶⁰ represents H or a suitable substituent.

Compounds of formula (I) wherein R⁸ is C₁₋₆alkyl substituted by thioxotriazolyl may be prepared from compounds of formula (I) wherein R⁸ is C₁₋₆alkyl substituted by CONHNH₂ by reaction with a compound of formula R⁶¹NCS, wherein R⁶¹ represents H or a suitable substituent such as C₁₋₆alkyl, in the presence of a base.

Suitable bases of use in the reaction include organic bases such as, for example, 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU). The reaction is conveniently effected in a suitable organic solvent, such as alcohol, e.g. butanol.

Compounds of formula (I) wherein R⁸ is C₁₋₆alkyl substituted by unsubstituted or substituted triazolyl may be prepared from compounds of formula (V)



wherein Hal is as previously defined, m is 1, 2, 3, 4, 5 or 6 and R⁶² is H or a group suitable as a substituent of the triazole ring, or convertible to such a group under the reaction conditions, in the presence of a base.

Suitable bases of use in the reaction include alkali metal carbonates, such as, for example, potassium carbonate.

Suitably R⁶² represents H, OCH₃ (which is converted to an oxo substituent under the reaction conditions) or CONH₂.

- 26 -

The reaction is conveniently effected in an anhydrous organic solvent, such as, for example, anhydrous dimethylformamide, preferably at elevated temperature, such as about 60°C.

Compounds of formula (I) wherein R^8 represents C_{1-6} alkyl substituted by $CONR^aC_{1-6}alkylR^{12}$ or $CONR^aheteroaryl$ may be prepared from compounds of formula (I) wherein R^8 is C_{1-6} alkyl substituted by CO_2H by reaction with an amine of formula $HNR^aC_{1-6}alkylR^{12}$ or $HNR^aheteroaryl$.

Alternatively, compounds of formula (I) in which R^7 represents, in particular, CO_2R^{16} may be converted into other compounds of formula (I). Thus compounds of formula (I) where R^7 is CO_2R^{16} may be reacted with a Grignard reagent of formula R^CMgHal where R^C is an alkyl group and Hal is as previously defined to give compounds where R^7 is a tertiary alcohol. Secondary alcohols may be prepared firstly by reduction of the ester moiety to an aldehyde using, for example, diisobutylaluminium hydride, followed by reaction with either R^CLi or R^CMgHal .

The aldehyde may also be used as a precursor for alkenyl intermediates where the group at position R^7 has the formula $-CH=CHR^d$, where R^d is $(CH_2)_sNR^9R^{10}$, $(CH_2)_sCO_2R^{16}$, $(CH_2)_sCONR^9R^{10}$ or $(CH_2)_sNR^9COR^{16}$ (where s is 0, 1 or 2 and R^9 , R^{10} and R^{16} are as previously defined). These compounds may be prepared by a Wittig reaction using, for example, $Ph_3P=CHR^d$ or $(EtO)_3P(O)=CHR^d$. These alkenyl intermediates may be reduced using, for example, catalytic hydrogenation to give compounds wherein R^7 is $(CH_2)_pNR^9R^{10}$, $(CH_2)_pCO_2R^{16}$, $(CH_2)_pCONR^9R^{10}$ or $(CH_2)_pNR^9COR^{16}$ and p is 2 to 4.

Compounds wherein R^7 is $CONR^9R^{10}$ may be prepared by the reaction of a compound of formula (I) wherein R^7 is CO_2R^{16} , and R^{16} is methyl, with an amine of the formula HNR^9R^{10} by known methods. Subsequent reduction using, for example, borane in tetrahydrofuran may be used to give a compound wherein R^7 is $(CH_2)_pNR^9R^{10}$ in which p is 1.

- 27 -


Where one or both of R^9 and R^{10} in the resultant amine is a hydrogen atom, the amine may be further converted into a compound wherein R^7 is $(CH_2)_pNR^9COR^{16}$ by reaction with, for example, an acyl chloride of the formula $R^{16}COCl$ by known methods.

5 Compounds wherein R^7 is CH_2OH may be prepared by the reduction of a compound of formula (I) wherein R^7 is CO_2R^{16} , and R^{16} is methyl, using, for example, lithium aluminium hydride. The primary alcohol may be used to prepare a compound of formula (I) wherein R^7 is $(CH_2)_pOR^{16}$ and p is 1 by reaction with a halide of the formula $R^{16}Hal$,
 10 where Hal is as previously defined in the presence of a suitable base such as sodium hydride.

 If a compound of the formula (I) is required in which R^6 is an alkyl group it may be prepared via a corresponding compound of the formula (I) in which R^7 is CO_2R^{16} group by reaction with KHMDS and an alkyl
 15 iodide, followed, if desired, by conversion of the CO_2R^{16} group as described above.

 According to a further general process (D), compounds of formula (I) wherein R^8 represents H and n is 2 or 3 may be prepared from a compound of formula (VI)

20




wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , and R^7 are as defined for formula (I) and x is 1 or 2 by reduction. Suitable reducing agents will be readily apparent to
 25 one skilled in the art and include, for example, borane or metallic hydrides,

- 28 -

such as lithium aluminium hydride or sodium borohydride. Borane is preferred.

Compounds of formula (VI) wherein R^6 is C_{1-6} alkyl and R^7 is H may be prepared by reductive amination of a compound of formula (VII)



wherein R^2 , R^3 , R^4 , and R^5 are as defined for formula (I) except that any reactive moiety is protected by a suitable protecting group and x is 1 or 2, using, for example, sodium cyanoborohydride or sodium triacetoxymethylborohydride and a compound of the formula R^1COR^6 where R^6 is C_{1-6} alkyl. The reaction is generally effected in a polar solvent such as acetic acid or methanol at a temperature between 0° and 50°C , conveniently at room temperature.

15 Alternatively, compounds of formula (VI) may be prepared by the reaction of a compound of formula (VII) with a compound of formula (III) in which R^{31} is a leaving group such as a halogen atom, for example, a bromine atom. The reaction is effected in the presence of a base, for example, potassium carbonate and in a suitable solvent such as dimethylformamide.

20 It will be appreciated that the product of the reductive amination method described herein will be a mixture of stereoisomers at the position of the group R^6 . For the subsequent preparation of a specific isomer of a compound of formula (I) wherein R^6 is C_{1-6} alkyl and R^7 is H, the mixture of stereoisomers may be resolved by conventional methods, for example, by column chromatography.

- 29 -

Methods for the preparation of intermediates of formula (VII) and formula (II) when R^{30} is NHR^3 are described, for example, in European Patent Specification No. 0 436 334.

Where they are not commercially available, the intermediates of formula (III) above may be prepared by the procedures described in the accompanying Examples or by alternative procedures which will be readily apparent to one skilled in the art.

Where the above-described processes for the preparation of the compounds according to the invention give rise to mixtures of stereoisomers these isomers may, if desired, be separated, suitably by conventional techniques such as preparative chromatography.

The novel compounds may be prepared in racemic form, or individual enantiomers may be prepared either by enantiospecific synthesis or by resolution. For example, compounds which contain a hydroxy group may be resolved into their component enantiomers by standard techniques, such as the formation of diastereomeric esters or amides, followed by chromatographic separation or separation by fractional crystallization and removal of the chiral auxiliary. Where they are intermediates, diastereomeric alcohols can then be used to prepare optically pure compounds of formula (I).

During any of the above synthetic sequences it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in *Protective Groups in Organic Chemistry*, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene and P.G.M. Wuts, *Protective Groups in Organic Synthesis*, John Wiley & Sons, 1991. The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

- 30 -

The exemplified compounds of this invention were tested by the methods set out at pages 36 to 39 of International Patent Specification No. WO 93/01165. The compounds were found to be active with IC₅₀ at the NK1 receptor of less than 150nM.

5 The compounds of this invention may be formulated as specifically illustrated at pages 35 to 36 of International Patent Specification No. WO 93/01165.

 The following Examples illustrate the preparation of compounds according to the invention.

10

Intermediate 1

cis-5-((3,5-Bis(trifluoromethyl))(α -methyl)benzylamino)-6-(4-fluorophenyl)-2-oxo-piperidine (isomers A and B)

15 5-Amino-6-(4-fluorophenyl)-2-oxo-piperidine (1.7g), citric acid (0.8g) and bis(trifluoromethyl)acetophenone (2.3g) were dissolved in methanol (30ml), powdered 4Å molecular sieves (2g) were added. After stirring for 5 minutes sodium cyanoborohydride (0.56g) was added, after stirring for a further 60 minutes the mixture was evaporated to dryness. The residue was suspended
20 in dichloromethane (30ml) and filtered through celite, the filtrate was then washed with 5% aqueous sodium bicarbonate (20ml), then dried (MgSO₄) and the solvent evaporated. The residue was purified by chromatography on silica (ethyl acetate:methanol, 9:1), to afford (isomer A) cis-5-((3,5-bis(trifluoromethyl))(α -methyl)benzylamino)-6-(4-fluorophenyl)-2-oxo-
25 piperidine (0.52g) mp 138°C, and, (isomer B) cis-5-((3,5-bis(trifluoromethyl))(α -methyl)benzylamino)-6-(4-fluorophenyl)-2-oxo-piperidine (0.21g) mp 120°C.

- 31 -

Intermediate 2cis-5-((3,5-Bis(trifluoromethyl))benzylamino)-6-(4-fluorophenyl)-2-oxo-piperidine

- 5 5-Amino-6-(4-fluorophenyl)-2-oxo-piperidine (1g), citric acid (0.5g) and bis(trifluoromethyl)benzaldehyde (1.3g) were dissolved in methanol (20ml), powdered 4Å molecular sieves (2g) were added. After stirring for 5 minutes sodium cyanoborohydride (0.34g) was added, after stirring for a further 60 minutes the mixture was evaporated to dryness. The residue was suspended
10 in dichloromethane (30ml) and filtered through celite, the filtrate was then washed with 5% aqueous sodium bicarbonate (20ml), then dried (MgSO₄) and the solvent evaporated. The residue was purified by chromatography on silica (ethyl acetate:methanol, 9:1), to afford cis-5-((3,5-bis(trifluoromethyl))benzylamino)-6-(4-fluorophenyl)-2-oxo-piperidine (1.1g)
15 ¹H NMR 1.64-1.72 (1H,m), 2.13-2.49 (3H,m), 2.95 (1H,br s), 3.77-3.86 (2H,m), 4.66 (1H,br s), 7.14 (2H,t,J=6Hz), 7.28(2H,tJ=6Hz), 7.76 (1H,s), 7.82 (2H,s), 7.88 (1H,s).

20 Intermediate 3

cis-3-((3,5-Bis(trifluoromethyl))benzylamino)-2-(4-fluorophenyl)piperidine

- cis-5-((3,5-Bis(trifluoromethyl))benzylamino)-6-(4-fluorophenyl)-2-oxo-piperidine (1g) was dissolved in tetrahydrofuran (15ml), a solution of borane
25 in THF was added dropwise (4.4ml of 1M solution), the resulting solution was then heated at reflux for 2 hours. The mixture was then cooled to room temperature and methanol (5ml) added dropwise, the resulting mixture was evaporated to dryness. The residue was dissolved in ethanol (20ml) potassium carbonate (1g) added and the mixture heated at reflux for 4 hours.

- 32 -

The mixture was then evaporated to dryness and the residue extracted with hot dichloromethane (2 x 20ml), ethereal HCl was added to the extracts and the product isolated by filtration. Recrystallisation from ethano/ether afforded *cis*-3-((3,5-bis(trifluoromethyl))benzylamino)-2-(4-fluorophenyl)piperidine as a dihydrochloride salt m.p. 290°C.

Example 1

10 *cis*-3-((3,5-Bis(trifluoromethyl))(α -methyl)benzylamino)-2-(4-fluorophenyl)piperidine (Isomer B)

Cis-5-((3,5-Bis(trifluoromethyl))(α -methyl)benzylamino)-6-(4-fluorophenyl)-2-oxo-piperidine (isomer B) (0.2g) was dissolved in tetrahydrofuran (15ml), a solution of borane in THF was added dropwise (2 ml of 1M soln), the resulting solution was then heated at reflux for 2 hours. The mixture was then cooled to room temperature and methanol (5 ml) added dropwise, the resulting mixture was evaporated to dryness. The residue was dissolved in ethanol (20ml) potassium carbonate (1g) added and the mixture heated at reflux for 4 hours. The mixture was then evaporated to dryness and the residue extracted with hot dichloromethane (2 x 20 ml), ethereal HCl was added to the extracts and the product isolated by filtration. Recrystallisation from ethanol/ether afforded *cis*-3-((3,5-bis(trifluoromethyl))(α -methyl)benzylamino)-2-(4-fluorophenyl)piperidine (Isomer B) as a dihydrochloride salt. m.p. 235°C

25

- 33 -

Example 2*cis*-5-[3-((3,5-Bis(trifluoromethyl))benzylamino)-2-(4-fluorophenyl)piperidin-1-ylmethyl]-2,4-dihydro-[1,2,4]-triazol-3-one

- 5 *cis*-3-((3,5-Bis(trifluoromethyl))benzylamino)-2-(4-fluorophenyl)piperidine dihydrochloride salt (0.68g) was partitioned between chloroform (50ml) and 10% aqueous potassium bicarbonate (20ml), the organic phase was separated, dried (Na₂SO₄) and evaporated to afford the free base (0.5g). The free base was dissolved in DMF (19ml) and K₂CO₃ (0.8g),
- 10 N-carbomethoxy-2-chloroacetamidrazone (0.2g) was added and the mixture heated to 60°C for 30 minutes. The mixture was then heated to 140°C for a further 2 hours, cooled to room temperature and the solvent removed at reduced pressure. The residue was purified by chromatography (ethyl acetate:methanol, 9:1), the oil isolated was dissolved in dichloromethane and
- 15 treated with ethereal HCl to afford the dihydrochloride salt, mp 225°C.

Example 3

- 20 *cis*-(±)-3-(2-Hydroxy-1-(4-fluorophenyl)ethylamino)-2-phenylpiperidine
- a) To a methanol (5ml) solution of (±) 5-amino-6-phenylpiperidin-2-one (1.67g, *cis:trans* isomers (6:1)) was added a solution of toluene-4-sulphonic acid monohydrate (1.8g) in methanol (5ml). The solid which formed by crystallization was removed by filtration to give pure *cis* isomer; mp 242-
- 25 244°C. A sample of this material (0.9g) was dissolved in warm water (10ml) and this solution was applied to a column containing Dowex™ 50W-X8 (H⁺ form, 20ml). After washing the column with water the product was eluted with dilute aqueous ammonia (2M) and the fractions containing product were evaporated to dryness. To a solution of the residue (0.32g) in

- 34 -

dimethylformamide (5ml) was added K_2CO_3 (0.7g) and 2-bromo-2-(4-fluorophenyl)acetic acid (0.42g). After stirring at room temperature for 16h, water (20ml) and ethyl acetate (20ml) were added and the organic phase washed with water, saturated in brine and dried ($MgSO_4$). The solvent was removed *in vacuo* and the residue purified on silica gel eluting with ethyl acetate to give, as a 0.7:1 mixture of diastereomers, methyl *cis* (\pm) 5-(N-(4-fluorophenylglycinate))-6-phenylpiperidin-2-one. 1H NMR (250MHz, $CDCl_3$) δ 7.45-7.09 (9H, m), 5.97 (0.4H, br d), 5.77 (br s), 4.90 (m), 4.78 (d, $J=3.6Hz$), 4.68 (d, $J=3.5Hz$), 4.6 (m), 4.05 (s+s), 3.61 (s, OMe), 3.54 (s, OMe), 3.08 (m), 2.87 (m), 2.75-2.3 (m), 1.90 (m). MS m/z (Cl^+) 357 (M+H).

b) A solution of the diastereomeric mixture (Example 3a, 0.32g) in tetrahydrofuran (4ml) was added to a solution of lithium aluminium hydride (1M in tetrahydrofuran, 2ml). The solution was stirred at room temperature for 15 minutes then at 40°C for 1 hour. Water (20ml) was added carefully to the solution followed by ethyl acetate (20ml). The organic phase was washed with saturated brine and dried ($MgSO_4$). The residue after evaporation was chromatographed on silica gel eluting with a gradient between CH_2Cl_2 and methanol, CH_2Cl_2 , aqueous ammonia (10:90:0.4) to give after crystallization from diethylether:

diastereomer A: mp=157-159°C, 1H NMR (250MHz, $CDCl_3$) δ 7.33-7.18 (5H, m, phenyl), 6.7 (dd, 2H, $J=8.72Hz$), 6.51 (2H, dd, $J=8.64Hz$, 5.44Hz, aryl), 3.89 (1H, d, $J=2.4Hz$, $NHCHPh$), 3.5 (2H, m), 3.22 (2H, m), 3.00 (1H, vbr s), 2.80 (1H, td, $J=12.01$, 2.75Hz), 2.70 (1H, d, $J=2.49Hz$), 1.90 (2H, m), 1.54 (2H, m) m/z (Cl^+) 315 (M+H); and

diastereomer B: mp=116-120°C, 1H NMR (250MHz, $CDCl_3$) δ 7.38-7.25 (5H, m), 7.0 (2H, dd, $J=6.6$, 8.7Hz), 6.74 (2H, dd, $J=8.7$, 8.7Hz), 3.8 (1H, d, $J=2.3Hz$), 3.28 (1H, dd, $J=10.6$, 4.5Hz), 3.18 (2H, m), 3.04 (1H, m), 2.78 (2H, m), 1.60 (3H, m), 1.37 (1H, m) m/z (Cl^+) 315 (M+H).

- 35 -


Example 4**5 cis-(±)-3-(2-Hydroxy-1-phenylethylamino)-phenylpiperidine**

The title compound was prepared in an analogous manner to that described in Example 3 using (±) 5-amino-6-phenylpiperidin-2-one and 2-bromo-2-phenylacetic acid as starting materials. MS m/z (Cl⁺) 297 (M+H).

- 36 -

CLAIMS:

1. A compound of formula (I), or a pharmaceutically acceptable salt or prodrug thereof:



wherein

- 10 n is 1, 2 or 3 and where any carbon atom of $(CH_2)_n$ may be substituted by R^4 and/or R^5 ;

- R^1 represents $(CH_2)_q$ phenyl, wherein q is zero, 1, 2 or 3, which may be optionally substituted in the phenyl ring by 1, 2 or 3 groups selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halo, cyano, nitro, trifluoromethyl, trimethylsilyl, $-OR^a$, $-SR^a$, $-SOR^a$, $-SO_2R^a$, $-NR^aR^b$, $-NR^aCOR^b$,
 15 $-NR^aCO_2R^b$, $-COR^a$, $-CO_2R^a$ and $-CONR^aR^b$;

- R^2 represents aryl selected from phenyl and naphthyl; heteroaryl selected from indazolyl, thienyl, furyl, pyridyl, thiazolyl, tetrazolyl and quinolyl; benzhydryl; or benzyl; wherein each aryl or heteroaryl moiety may be substituted by C_{1-6} alkyl, C_{1-6} alkoxy, halo or trifluoromethyl;

- 20 R^3 represents H or C_{1-6} alkyl;

- R^4 and R^5 each independently represent H, halo, C_{1-6} alkyl, oxo, CO_2R^a or $CONR^aR^b$;

- R^6 represents H or C_{1-6} alkyl;

- 37 -

R⁷ represents H, C₁₋₆alkyl optionally substituted by a hydroxy group, or (CH₂)_pNR⁹R¹⁰, CO₂R¹⁶, CONR⁹R¹⁰, (CH₂)_pCO₂R¹⁶, (CH₂)_pCONR⁹R¹⁰, (CH₂)_pNR⁹COR¹⁶, (CH₂)_pNHSO₂R¹¹, (CH₂)_pOR¹⁶, (CH₂)_pOC(O)R⁹, (CH₂)_pOC₁₋₄alkylCOR¹⁷, or phenyl optionally substituted
 5 by 1, 2 or 3 groups selected from C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, halo, cyano, nitro, trifluoromethyl, trimethylsilyl, -OR^a, -SR^a, -SOR^a, -SO₂R^a, -NR^aR^b, -NR^aCOR^b, -NR^aCO₂R^b, -COR^a, -CO₂R^a and -CONR^aR^b;

R⁸ represents H, COR^a, CO₂R^a, COCONR^aR^b, COCO₂R^a, C₁₋₆alkyl optionally substituted by a group selected from (CO₂R^a,
 10 CONR^aR^b, hydroxy, cyano, COR^a, NR^aR^b, C(NO₂)NR^aR^b, CONHphenyl(C₁₋₄alkyl), COCO₂R^a, CONHN^aR^b, C(S)NR^aR^b, CONR^aC₁₋₆alkylR¹², CONR¹³C₂₋₆alkynyl, CONR¹³C₂₋₆alkenyl, COCONR^aR^b, CONR^aC(NR^b)NR^aR^b, CONR^aheteroaryl, and phenyl optionally substituted by one or more substituents selected from C₁₋₆alkyl,
 15 C₁₋₆alkoxy, halo and trifluoromethyl) or C₁₋₆alkyl, optionally substituted by oxo, substituted by an optionally substituted aromatic heterocycle;

with the proviso that when R⁶ and R⁷ both represent H, R⁸ represents C₁₋₆alkyl optionally substituted by oxo, substituted by an optionally substituted aromatic heterocycle selected from pyrrolyl, pyrazolyl,
 20 pyrazinyl, pyridazinyl, oxadiazolyl, thiadiazolyl, imidazolyl, benzimidazolyl, benzoxazolyl, benzthiophenyl, benzofuranyl and indolyl, or a substituted aromatic heterocycle selected from thienyl, furyl, pyridyl, triazolyl, tetrazolyl, thiazolyl, oxazolyl, quinolyl, isoxazolyl and isothiazolyl;

R^a and R^b each independently represent H, C₁₋₆alkyl, trifluoromethyl or phenyl optionally substituted by C₁₋₆alkyl, halo or trifluoromethyl;
 25

R⁹ and R¹⁰ each independently represent H or C₁₋₆alkyl;

R¹¹ represents NR¹⁴R¹⁵ or an optionally substituted aromatic or non-aromatic azacyclic or azabicyclic group;

- 38 -

R¹² represents OR^a, CONR^aR^b or heteroaryl;

R¹³ represents H or C₁₋₆alkyl;


R¹⁴ and R¹⁵ each independently represent H, C₁₋₆alkyl, phenyl optionally substituted by one or more of C₁₋₆alkyl, C₁₋₆alkoxy, halo or trifluoromethyl, or phenylC₁₋₄alkyl optionally substituted in the phenyl ring by one or more of C₁₋₆alkyl, C₁₋₆alkoxy, halo or trifluoromethyl;

R¹⁶ represents C₁₋₆alkyl;

R¹⁷ represents C₁₋₆alkoxy, amino, C₁₋₄alkylamino or di(C₁₋₄alkyl)amino; and

p is 1 to 4.

2. A compound as claimed in claim 1 of formula (Ia), or a pharmaceutically acceptable salt or prodrug thereof:



wherein

R¹, R², R³, R⁴, R⁵, R⁶, R⁸ and n are as defined in claim 1;


and

R⁷ represents C₁₋₆alkyl optionally substituted by a hydroxy group, or (CH₂)_pNR⁹R¹⁰, CO₂R¹⁶, CONR⁹R¹⁰, (CH₂)_pCO₂R¹⁶, (CH₂)_pCONR⁹R¹⁰, (CH₂)_pNR⁹COR¹⁶, (CH₂)_pNHSO₂R¹¹, (CH₂)_pOR¹⁶, (CH₂)_pOC(O)R⁹, (CH₂)_pOC₁₋₄alkylCOR¹⁷, or phenyl optionally substituted by 1, 2 or 3 groups selected from C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, halo,

- 39 -

cyano, nitro, trifluoromethyl, trimethylsilyl, $-OR^a$, $-SR^a$, $-SOR^a$, $-SO_2R^a$, $-NR^aR^b$, $-NR^aCOR^b$, $-NR^aCO_2R^b$, $-COR^a$, $-CO_2R^a$ and $-CONR^aR^b$.

3. A compound as claimed in claim 1 or claim 2 of formula
5 (Ib), or a pharmaceutically acceptable salt or prodrug thereof:




wherein

10 R^2 , R^3 , R^4 , R^5 , R^6 , R^8 and n are as defined in claim 1;
 R^{18} represents C_{1-6} alkyl optionally substituted by a hydroxy
 group; and
 R^{19} represents phenyl optionally substituted by 1, 2 or 3
 groups selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halo, cyano, nitro,
 15 trifluoromethyl, trimethylsilyl, $-OR^a$, $-SR^a$, $-SOR^a$, $-SO_2R^a$, $-NR^aR^b$,
 $-NR^aCOR^b$, $-NR^aCO_2R^b$, $-COR^a$, $-CO_2R^a$ and $-CONR^aR^b$.

4. A compound as claimed in any one of claims 1 to 3 of
 formula (Ic), or a pharmaceutically acceptable salt or prodrug thereof:

20

- 40 -



wherein

R^3 and R^8 are as defined in claim 1;

5 R^{20} and R^{21} independently represent H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, chloro, bromo, fluoro, iodo, cyano, nitro, trifluoromethyl, trimethylsilyl, OR^a , SR^a , SOR^a , SO_2R^a , NR^aR^b , NR^aCOR^b , $NR^aCO_2R^b$, COR^a , CO_2R^a or $CONR^aR^b$, where R^a and R^b are as previously defined;

10 R^{22} is methyl optionally substituted by hydroxy;


R^{23} is H or methyl; and

R^{24} represents phenyl or benzhydryl wherein any of the phenyl rings of the phenyl or benzhydryl moieties may optionally be substituted by halo or trifluoromethyl.

15

5. A compound as claimed in claim 1 of formula (Id), or a pharmaceutically acceptable salt or prodrug thereof:

- 41 -




wherein

R¹, R², R³, R⁴, R⁵, and n are as defined in claim 1; and

5 R⁸ represents C₁₋₆alkyl, optionally substituted by oxo,
substituted by an optionally substituted aromatic heterocycle selected from
pyrrolyl, pyrazolyl, pyrazinyl, pyridazinyl, oxadiazolyl, thiadiazolyl, imidazolyl,
benzimidazolyl, benzoxazolyl, benzothiophenyl, benzofuranyl and indolyl, or a
substituted aromatic heterocycle selected from thienyl, furyl, pyridyl, triazolyl,
10 tetrazolyl, thiazolyl, oxazolyl, quinolyl, isoxazolyl and isothiazolyl.

6. A compound as claimed in claim 1 or claim 5 of formula
(Ie), or a pharmaceutically acceptable salt or prodrug thereof.



15


wherein R², R³, R⁴, R⁵, R⁸ and n are as defined in claim 1; and

R¹⁹ represents phenyl optionally substituted by 1, 2 or 3
groups selected from C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, halo, cyano, nitro,

- 42 -

trifluoromethyl, trimethylsilyl, $-OR^a$, $-SR^a$, $-SOR^a$, $-SO_2R^a$, $-NR^aR^b$, $-NR^aCOR^b$, $-NR^aCO_2R^b$, $-COR^a$, $-CO_2R^a$ and $-CONR^aR^b$.

7. A compound as claimed in any one of claims 1, 5 or 6 of formula (If), or a pharmaceutically acceptable salt or prodrug thereof:



wherein

- 10 R^3 and R^8 are as defined in claim 1;
 R^{20} and R^{21} independently represent H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, chloro, bromo, fluoro, iodo, cyano, nitro, trifluoromethyl, trimethylsilyl, OR^a , SR^a , SOR^a , SO_2R^a , NR^aR^b , NR^aCOR^b , $NR^aCO_2R^b$, COR^a , CO_2R^a or $CONR^aR^b$, where R^a and R^b are as
 15 previously defined; and

R^{24} represents phenyl or benzhydryl wherein any of the phenyl rings of the phenyl or benzhydryl moieties may optionally be substituted by halo or trifluoromethyl.

- 20 8. A compound selected from:
cis-3-((3,5-bis(trifluoromethyl))(α -methyl)benzylamino)-2-(4-fluorophenyl) piperidine;

- 43 -

cis-[3-((3,5-bis(trifluoromethyl))benzylamino)-2-(4-fluorophenyl)piperidine-1-ylmethyl]-2,4-dihydro-[1,2,4]-triazol-3-one;
or a pharmaceutically acceptable salt or prodrug thereof.

5 9. A compound as claimed in any preceding claim for use in therapy.

 10. A pharmaceutical composition comprising a compound as claimed in any one of Claims 1 to 8 in association with a pharmaceutically acceptable carrier or excipient.

10

 11. A method for the treatment or prevention of physiological disorders associated with an excess of tachykinins, which method comprises administration to a patient in need thereof of a tachykinin reducing amount of a compound according to Claim 1, or a pharmaceutically acceptable salt thereof, or a composition comprising a compound according to Claim 1, or a pharmaceutically acceptable salt thereof.

15

 12. A method according to Claim 11 for the treatment or prevention of pain or inflammation.

20

 13. A method according to Claim 11 for the treatment or prevention of migraine.

 14. A method according to Claim 11 for the treatment or prevention of emesis.

25

- 44 -

15. The use of a compound as claimed in Claim 1 for the manufacture of a medicament for the treatment of a physiological disorder associated with an excess of tachykinins.

5 16. The use of a compound as claimed in claim 1 for the manufacture of a medicament for the treatment of pain or inflammation.

17. The use of a compound as claimed in Claim 1 for the manufacture of a medicament for the treatment of migraine.


10

18. The use of a compound as claimed in Claim 1 for the manufacture of a medicament for the treatment of emesis.

19. A process for the preparation of a compound of formula (I) which comprises:

15

(A) reacting a compound of formula (II) with a compound of formula (III):



20

wherein R¹, R², R⁴, R⁵, R⁶, R⁷, R⁸ and n are as defined in claim 1, or a protected derivative thereof, and one of R³⁰ and R³¹ represents a leaving

- 45 -


group and the other of R^{30} and R^{31} represents NHR^3 , where R^3 is as defined in claim 1; in the presence of a base; or

(B) wherein R^8 is hydrogen, by the reductive amination of a compound of formula (II) in which R^{30} is the group NHR^3 , with a compound of formula R^1COR^7 , in the presence of a reducing agent; or

(C) by interconversion of a compound of formula (I) into another compound of formula (I); or

10

(D) wherein R^8 represents H and n is 2 or 3, by reduction of a compound of formula (VI)



15

wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , and R^7 are as defined in claim 1 and x is 1 or 2;

each process being followed, where necessary, by the removal of any protecting group where present;

20

and when the compound of formula (I) is obtained as a mixture of enantiomers or diastereoisomers, optionally resolving the mixture to obtain the desired enantiomer;

and/or, if desired, converting the resultant compound of formula (I) or a salt thereof into a pharmaceutically acceptable salt thereof.

25

INTERNATIONAL SEARCH REPORT

Intern: J Application No
PCT/GB 95/00153

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D211/56 C07D401/06 A61K31/445

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO,A,93 01170 (PFIZER) 21 January 1993 see page 33 - page 34; claim 1 ---	1,15
Y,P	CHEMICAL ABSTRACTS, vol. 120, no. 5, 31 January 1994 Columbus, Ohio, US; abstract no. 45616t, page 74; cited in the application see abstract	1,15
Y	& EUR. J. PHARMACOL., vol. 250, no.1, 1993 pages r5-r6, TATTERSALL,F;RYCROFT,W.;HARGREAVES,R.;HILL ,R 'The tachykinin NK1 receptor antagonist CP-99994 attenuates cisplatin induced emesis in the ferret' --- -/--	1,15

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

* & * document member of the same patent family

Date of the actual completion of the international search

10 April 1995

Date of mailing of the international search report

- 9. 05. 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Lauro, P

INTERNATIONAL SEARCH REPORT

Intern. d Application No
PCT/GB 95/00153

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP,A,0 436 334 (PFIZER) 10 July 1991 see page 17, line 50 - page 19, line 22; example 84 ---	1,15
P,A	WO,A,94 03445 (PFIZER) 17 February 1994 see examples 4,8 ---	1
A	WO,A,93 21181 (MERCK SHARP & DOHME) 28 October 1993 see page 15, line 24 - page 18, line 28; claim 1 -----	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Application No

PCT/GB 95/00153

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
WO-A-9301170	21-01-93	US-A-	5232929	03-08-93
		CA-A-	2111461	21-01-93
		EP-A-	0594636	04-05-94
		JP-T-	6508828	06-10-94
		US-A-	5332817	26-07-94

EP-A-436334	10-07-91	WO-A-	9109844	11-07-91
		AT-T-	115127	15-12-94
		CN-A-	1087083	25-05-94
		DE-D-	69014848	19-01-95
		EP-A-	0558156	01-09-93
		ES-T-	2064667	01-02-95
		PL-B-	163967	30-06-94
		PL-B-	164203	30-06-94
		PL-B-	164204	30-06-94
PL-B-	164205	30-06-94		

WO-A-9403445	17-02-94	AU-B-	4396193	03-03-94
		CA-A-	2141051	17-02-94
		FI-A-	933455	05-02-94

WO-A-9321181	28-10-93	AU-B-	4076593	18-11-93
		CA-A-	2133077	28-10-93
		EP-A-	0636130	01-02-95
		AU-B-	4713993	14-02-94
		CA-A-	2138650	03-02-94
		WO-A-	9402461	03-02-94
